arXiv:1111.1608v1 [math.CO] 7 Nov 2011

EQUIPARTITIONS OF MEASURES BY TWO HYPERPLANES

RADE T. ŽIVALJEVIĆ

Abstract. We compute a primary cohomological obstruction to the existence of an equipartition for j mass distributions in $\mathbb{R}^d$ by two hyperplanes in the case $2d-3j=1$. The central new result is that such an equipartition always exists if $d=6\cdot 2^k+2$ and $j=4\cdot 2^k+1$ which for $k=0$ reduces to the main result of the paper P. Mani-Levitska et al., Topology and combinatorics of partitions of masses by hyperplanes, *Adv. Math.* 207 (2006), 266–296. The theorem follows from a Borsuk-Ulam type result claiming the non-existence of a $\mathbb{D}_8$-equivariant map $f: S^d\times S^d\to S(W^{\oplus j})$ for an associated real $\mathbb{D}_8$-module W. This is an example of a genuine combinatorial geometric result which involves $\mathbb{Z}/4$-torsion in an essential way and cannot be obtained by the application of either Stiefel-Whitney classes or cohomological index theories. The method opens a possibility of developing an "effective primary obstruction theory" based on G-manifold complexes, with applications in geometric combinatorics, discrete and computational geometry, and computational algebraic topology.

1. Introduction

1.1. **The equipartition problem.** An old problem in combinatorial geometry is to determine when j measurable sets in $\mathbb{R}^d$ admit an *equipartition* by a collection of k hyperplanes; if this is possible we say that the triple (d,j,k) is admissible. In this generality the problem was formulated by Grünbaum in [5] but the problem clearly stems from the classical results of Steinhaus [20] and Stone and Tukey [21] on the "ham sandwich theorem".

Among the highlights of the theory of hyperplane partitions of measurable sets (measures) are Hadwiger's equipartition of a single mass distribution in $\mathbb{R}^3$ by three hyperplanes [8], results of Ramos [16] who proved that $(5,1,4),(9,3,3),(9,5,2)$ are admissible triples, and a result of Mani-Levitska et al. [13] who established a 2-equipartition for 5 measures in $\mathbb{R}^8$. More recent is the proof of the existence of a 4-equipartition in $\mathbb{R}^4$ for measures that admit some additional symmetries (Živaljević [26]) and a result of Matschke [15] describing a general reduction procedure for admissible triples.

For an account of known results, history of the problem, and an exposition of equivariant topological methods used for its solution, the reader is referred to [13]. The landmark paper of E. Ramos [16] is a valuable source of information and a link with earlier results on the discrete and computational geometry of half-space range queries (D. Dobkin, H. Edelsbrunner, M. Paterson, A. Yao, F. Yao). The chapter "Topological Methods" in CRC Handbook of Discrete and Computational

---

Date: October 30, 2011 (first version, April 10, 2011).

Supported by the Serbian Ministry of Science and Technology, Grants 174020 and 174034 .

Geometry [25], provides an overview of the general *configuration space/test map*-scheme for applying equivariant topological methods on problems od geometric combinatorics and discrete and computational geometry.

Definition 1.1. Suppose that $\mathcal{M} = \{\mu_1, \mu_2, \dots, \mu_j\}$ is a collection of j *continuous mass distributions* (measures) defined in $\mathbb{R}^d$, meaning that each μ_j is a finite, positive, σ-additive Borel measure, absolutely continuous with respect the Lebesgue measure. If $\mathcal{H} = \{H_i\}_{i=1}^k$ is a collection of k hyperplanes in $\mathbb{R}^d$ in general position, the connected components of the complement $\mathbb{R}^d \setminus \cup\, \mathcal{H}$ are called (open) k-orthants. The definition can be clearly extended to the case of degenerate collections $\mathcal{H}$, when some of the k-orthants are allowed to be empty. A collection $\mathcal{H}$ is an *equipartition*, or more precisely a k-equipartition for $\mathcal{M}$ if

$$\mu_i(O) = \mu_i(\overline{O}) = \frac{1}{2^k}\mu_i(\mathbb{R}^d)$$

for each of the measures $\mu_i \in \mathcal{M}$ and for each k-orthant O associated to $\mathcal{H}$.

A triple (d, j, k) is called *admissible* if for each collection $\mathcal{M}$ of j continuous measures on $\mathbb{R}^d$ there exists an equipartition of $\mathcal{M}$ by k hyperplanes. It is not difficult to see that if (d, j, k) is admissible that $(d + 1, j, k)$ is also admissible[1] which motivates the following general problem.

Problem 1.2. The general measure equipartition problem is to determine or estimate the function

$$\Delta(j, k) := \min\{d \mid (d, j, k) \text{ is admissible}\}$$

or equivalently to find the minimum dimension d such that for each collection $\mathcal{M}$ of j continuous measures on $\mathbb{R}^d$, there exists an equipartition of $\mathcal{M}$ by k hyperplanes.

2. New results

Theorem 2.1 is our central new result about equipartitions of masses by two hyperplanes. For $k = 0$ it reduces to the result that the triple $(8, 5, 2)$ is admissible, which is the main result of [13]. The reader is referred to references [16, 13, 15] for an analysis explaining the special role of equipartition problems associated to the triples of the form $(d, j, 2)$ where $2d - 3j = 1$. We emphasize that the admissibility of all triples listed in Theorem 2.1 cannot be established either by the parity count results of Ramos [16], or by the use of both Stiefel-Whitney characteristic classes and the ideal-valued cohomology index theory with $\mathbb{Z}/2$-coefficients.

Theorem 2.1. *Each collection of* $j = 4 \cdot 2^k + 1$ *measures in* $\mathbb{R}^d$ *where* $d = 6 \cdot 2^k + 2$ *admits an equipartition by two hyperplanes. In light of the lower bound* $\Delta(j, 2) \geq 3j/2$*, established by Ramos in* [16]*, this result implies that for each integer* $k \geq 0$*,*

$$\Delta(4 \cdot 2^k + 1, 2) = 6 \cdot 2^k + 2. \tag{2.2}$$

Theorem 2.1 is deduced from the following Borsuk-Ulam type result about maps equivariant with respect to the dihedral group actions.

---

[1] More general reduction results for admissible triples can be found in [15] and [16].

Theorem 2.3. *There does not exist a* $\mathbb{D}_8$*-equivariant map*

$$f : S^d \times S^d \to S(W^{\oplus j}) \tag{2.4}$$

where $\mathbb{D}_8$ *is the dihedral group of order eight,* $d = 6 \cdot 2^k + 2, j = 4 \cdot 2^k + 1$ *for some integer* $k \geq 0$*, and* W *is the representation space of the real* 3*-dimensional* $\mathbb{D}_8$*-representation described in Section 4.2.*

Moreover, a first obstruction to the existence of (2.4) lies in the equivariant cohomology group $\mathcal{H}^{2d-1}_{\mathbb{D}_8}(S^d \times S^d, \mathcal{Z}) \cong \mathbb{Z}/4$*, described in Sections 7 and 8, where* $\mathcal{Z} = H_{2d-2}(S(W^{\oplus j}); \mathbb{Z})$*. The obstruction vanishes unless*

$$d = 6 \cdot 2^k + 2 \quad \text{and} \quad j = 4 \cdot 2^k + 1$$

when it turns out to be equal to $2X$ *where* X *is a generator of the group* $\mathbb{Z}/4$*.*

The fact that the obstruction $2X \in \mathbb{Z}/4$ is divisible by 2 is precisely the reason why Theorem 2.3 is not accessible by the methods based on $\mathbb{Z}/2$-coefficients (parity count [16], Stiefel-Whitney classes, $\mathbb{D}_8$-equivariant index theory with $\mathbb{Z}/2$-coefficients). As the elaborate spectral sequences calculations [1] demostrate, the methods of $\mathbb{D}_8$-equivariant index theory with $\mathbb{Z}$-coefficients are not sufficient either. This may be somewhat of an accident since, as shown in [3], the $\mathbb{Z}/4$-torsion is often present in related cohomology calculations.

Remark 2.5. In light of the reduction procedure of Matschke [15], who proved the inequality $\Delta(j, k) \leq \Delta(j+1, k) - 1$, it is interesting to test if Theorem 2.1 generates some new admissible triples aside from those implied by the inequality (2.2). The answer is no, since the inequality $\Delta(2^{k+1}, 2) \leq 3 \cdot 2^k$ was established already by Ramos in [16].

3. Related results and background information

Results about partitions of measures by hyperplanes have found numerous applications. The survey [25] covers the results from the first half of the past decade or older. For this reason we briefly review some of the most interesting recent developments illustrating the relevance and importance of hyperplane partitions for different areas of mathematics.

3.1. **Polyhedral partitions of measures.**

Equipartitions of measures by k-orthants (Definition 1.1) are a special case of equipartitions into polyhedral regions. Very interesting and far reaching results in this direction have been recently obtained by Soberon [18], Karasev [10], and Aronov and Hubard [9].

Referring the reader to original papers, we select only one of these results for illustrating a relationship between hyperplane partitions and polyhedral partitions of measures. The following theorem is attributed to Gromov [4], since it is an immediate consequence of the construction leading to the Gromov's Borsuk-Ulam theorem [4, Section 5]. However Karasev [10] was probably the first who recognized its power and potential for various generalizations and ramifications.

Definition 3.1. Let T_d be a binary tree of height d, with $2^d - 1$ internal nodes (root included) and 2^d external nodes (leaves). A "decorated binary three" $(T_d, \{H_\nu\}_{\nu \in N_d})$ has an oriented hyperplane H_ν (and the positive half-space H^+_ν) associated to each of the internal nodes ν of T_d. Given a decorated binary tree $(T_d, \{H_\nu\}_{\nu\in N_d})$ the associated *tree-like hyperplane partition* $\Pi = \{\Pi_\pi\}_{\pi\in L_d}$ of $\mathbb{R}^n$ is the decomposition $\mathbb{R}^n = \cup_{\pi\in L_d} \Pi_\pi$ into polyhedral regions $\Pi_\pi := \cap\{H_\nu^+\}_{\nu\prec\pi}$.

Theorem 3.2. *Let $\mu_1, \mu_2, \dots, \mu_n$ be a collection of n continuous mass distributions (Definition 1.1) on $\mathbb{R}^n$. Then there exists a tree-like hyperplane partition $\Pi = \{\Pi_\pi\}_{\pi\in L_d}$ of $\mathbb{R}^n$ which is an equipartition for each of the measures μ_j in the sense that*

$$\forall j, \forall \pi \in L_d,\ \mu_j(\Pi_\pi) = \frac{1}{2^d}\mu_j(\mathbb{R}^d).$$

3.2. **Polynomial measure partitions theorems.** Theorem 2.1, being a relative of the Ham Sandwich Theorem, has some standard consequences and extensions. One of them is the Polynomial Ham Sandwich Theorem [21], which has recently found striking applications to some old problems of discrete and computational geometry, see [7], [19] and the references in these papers. These breakthroughs have generated a lot of interest and enthusiasm, see the reviews of J. Matoušek [14], M. Sharir [17] and T. Tao [22].

Both theorems 2.1 and 3.2, as well other measure partitions results, including those listed in the review paper [25], have immediate polynomial versions. They can be obtained by the usual Veronese map $\mathbb{R}^d \hookrightarrow \mathbb{R}^D$, or some of its variations. It remains to be seen if some of these polynomial measure partition results can be used as a natural tool which can replace the standard polynomial ham sandwich theorem in some applications.

4. Preliminaries, definitions, notation

4.1. Manifold complexes.

Definition 4.1. A space X is called a *manifold pre-complex* if it is either a compact topological manifold (with or without boundary) or if it is obtained by gluing a compact topological manifold with boundary to a manifold pre-complex via a continuous map of the boundary.

This definition appeared in [11] (Chapter 9) where manifold pre-complexes are referred to as *nice spaces.* One can *mutatis mutandis* modify the Definition 4.1 by allowing different kinds of "manifolds". For example X is a pseudomanifold pre-complex (orientable manifold pre-complex, complex manifold pre-complex, etc.), if the constituent "manifolds" are pseudomanifolds (orientable manifolds, complex manifolds, etc.).

A manifold pre-complex should be seen as a straightforward generalization of a (finite) CW-complex. However a CW-complex has a natural filtration (and an associated rank function) so for this reason we slightly modify the definition and introduce *manifold complexes.*

Definition 4.2. A space X with a finite filtration

$$X_0 \subset X_1 \subset \ldots \subset X_{n-1} \subset X_n \tag{4.3}$$

is called a *manifold complex* (orientable manifold complex, etc.) if

(1) X_0 is a finite set of points (0-dimensional manifold);
(2) For each $k \leq n$, $X_k = X_{k-1} \cup_\phi Y_k$ where Y_k is a compact k-dimensional manifold with boundary and $\phi : \partial Y_k \to X_{k-1}$ is a continuous map.

As a variation on a theme we introduce manifold complexes with an action of a finite group G.

Definition 4.4. Let G be a finite group. A G-space X which is also a manifold complex in the sense of Definition 4.2 is called a G-manifold complex if G preserves the filtration (4.3).

4.2. **Dihedral group $\mathbb{D}_8$.** For basic notation and standard facts about group actions the reader is referred to [2]. A representation space V for a given G-representation $\rho : G \to GL(V)$ is also referred to as a (real or complex) G-module. $S(V)$ is the unit sphere in an orthogonal (or unitary) G-representation space V. $X * Y$ is the join of two spaces X and Y, and a standard fact is that if U and V are two orthogonal G-modules, $S(U \oplus V)$ and $S(U) * S(V)$ are isomorphic as G-spaces.

$\mathbb{D}_8$ is the dihedral group of order 8. $\Lambda = \mathbb{Z}[\mathbb{D}_8]$ is the integral group ring of $\mathbb{D}_8$ and Λx denotes the free, one-dimensional Λ-module generated by x. In this paper x is often a fundamental class of an orientable $\mathbb{D}_8$-pseudomanifold with boundary.

As the group of symmetries of the square $Q = \{(x, y) \in \mathbb{R}^2 \mid 0 \leq |x|, |y| \leq 1\}$ the dihedral group $\mathbb{D}_8$ has three distinguished involutions α, β and γ where

$$\alpha(x, y) = (-x, y), \quad \beta(x, y) = (x, -y), \quad \gamma(x, y) = (y, x). \tag{4.5}$$

A standard presentation of $\mathbb{D}_8$ is,

$$\mathbb{D}_8 = \langle \alpha, \beta, \gamma \mid \alpha^2 = \beta^2 = \gamma^2 = 1, \alpha\beta = \beta\alpha, \alpha\gamma = \gamma\beta, \beta\gamma = \gamma\alpha \rangle.$$

The real two-dimensional representation $\rho : \mathbb{D}_8 \to O(2)$ arising from the action on the square is denoted by U. Let $W := U \oplus \epsilon$ where ϵ is the one-dimensional (real) $\mathbb{D}_8$-representation, such that

$$\alpha \cdot x = -x, \quad \beta \cdot x = -x, \quad \gamma \cdot x = x. \tag{4.6}$$

Interpreting $\mathbb{D}_8$ is a Sylow 2-subgroup of the symmetric group S_4 we see that the $\mathbb{D}_8$-module W is isomorphic to the restriction of the reduced permutation representation of S_4 to the dihedral group.



FIGURE 1. $\mathbb{D}_8$-module W as a permutation representation.

More explicitly, as shown in Figure 1, the permutations associated to the basic involutions α, β and γ are:

$$\alpha = \begin{pmatrix} 0 & 1 & 2 & 3 \\ 1 & 0 & 3 & 2 \end{pmatrix}, \quad \beta = \begin{pmatrix} 0 & 1 & 2 & 3 \\ 3 & 2 & 1 & 0 \end{pmatrix}, \quad \gamma = \begin{pmatrix} 0 & 1 & 2 & 3 \\ 0 & 3 & 2 & 1 \end{pmatrix} \tag{4.7}$$

5. The topology of the equipartition problem

The problem of deciding if for a given collection $\{\mu_1, \dots, \mu_j\}$ of measures in $\mathbb{R}^d$ there exists an equipartition by an ordered pair (H_0, H_1) of (oriented) hyperplanes, can be reduced to a topological problem via the usual *Configuration Space/Test Map Scheme*, see [25, Section 14.1] or [13, Section 2.3]. Here is a brief outline of this construction.

The configuration space, or the space of all candidates for the equipartition, is the space of all ordered pairs (H_0, H_1) of oriented hyperplanes. After a suitable compactification this space can be identified as $S^d \times S^d$. The natural group of symmetries for the equipartition problem is the dihedral group $\mathbb{D}_8$ and the associated action on $S^d \times S^d$ is given by formulas (4.5).

The importance of the representation W stems from the fact that it naturally arises [13, Section 2.3.3] as the associated *Test Space* for a single (probabilistic) measure μ. Indeed for each ordered pair (H_0, H_1) of oriented hyperplanes there is an associated collection of hyperorthants O_0, O_1, O_2, O_3 (Figure 1 (a)). Then $\mu(O_\nu)$, $\nu = 0, 1, 2, 3$ are naturally interpreted as the *barycentric coordinates* of a point $v = \mu(O_0)v_0 + \mu(O_1)v_1 + \mu(O_2)v_2 + \mu(O_3)v_3 \in W$ (Figure 1 (b)) and the action of $\mathbb{D}_8$ on $S^d \times S^d$ induces an action on these barycentric coordinates which is precisely the action on W as a $\mathbb{D}_8$-module described in Section 4.2. Assume that the barycenter of the tetrahedron is the origin, i.e. $(1/4)(v_0 + v_1 + v_2 + v_3) = 0 \in W$. The map

$$F_\mu : S^d \times S^d \to W, \quad (H_0, H_1) \mapsto \mu(O_0)v_0 + \mu(O_1)v_1 + \mu(O_2)v_2 + \mu(O_3)v_3$$

has the property that (H_0, H_1) is an equipartition for μ if and only if (H_0, H_1) is a zero of F_μ. More generally $z := (H_0, H_1)$ is an equipartition for the collection $\{\mu_1, \dots, \mu_j\}$ of probability measures if and only if $z = (H_0, H_1)$ is a common zero of the associated test maps F_{μ_j},

$$F_{\mu_1}(z) = F_{\mu_2}(z) = \dots = F_{\mu_j}(z) = 0.$$

Summarizing we have the following proposition.

Proposition 5.1. *A triple $(d, j, 2)$ is admissible if each $\mathbb{D}_8$-equivariant map*

$$F : S^d \times S^d \to W^{\oplus j}$$

has a zero, or equivalently if there does not exist a $\mathbb{D}_8$-equivariant map

$$f : S^d \times S^d \to S(W^{\oplus j}). \tag{5.2}$$

6. STANDARD ADMISSIBLE FILTRATION OF $S^n \times S^n$

In order to prove the non-existence of an equivariant map (5.2) we apply the equivariant obstruction theory in the form outlined in Section 11. The first step is a construction of an appropriate filtration on $S^n \times S^n$ which is admissible in the sense of Definition 11.5. Less formally, we turn $S^n \times S^n$ into a $\mathbb{D}_8$-manifold complex in the sense of Definition 4.4 by allowing orientable $\mathbb{D}_8$-manifolds with corners and mild singularities.

For $i = 1, \ldots, n+1$ define $\pi_i : S^n \times S^n \to \mathbb{R}^2$, $\pi_i(x, y) := (x_i, y_i)$ as the restriction of the obvious projection map. The maps π_i are clearly $\mathbb{D}_8$-equivariant and the images $\mathrm{Image}(\pi_i) = \{(x_i, y_i) \mid -1 \leq x_i, y_i \leq +1\} =: Q_i$ are squares which are here referred to as "$\mathbb{D}_8$-screens", Figure 2. The screens Q_i admit a $\mathbb{D}_8$-invariant triangulation which is the starting point for the construction of an admissible filtration on $S^n \times S^n$.



FIGURE 2. $\mathbb{D}_8$-screens for $S^2 \times S^2 \subset \mathbb{R}^3 \times \mathbb{R}^3$.

The filtration can be informally described as follows. The manifold $S^n \times S^n$ is fibered over the first screen Q_1 with a generic fiber homeomorphic to $S^{n-1} \times S^{n-1}$. The fiber $S^{n-1} \times S^{n-1}$ itself is fibered over the second screen Q_2 with $S^{n-2} \times S^{n-2}$ as a generic fiber, etc. This priority order of screens together with their minimal $\mathbb{D}_8$-invariant triangulations are used to define a version of "lexicographic filtration" of $S^n \times S^n$. For the intended application of the obstruction theory methods from Section 11 it will be sufficient give a precise description only for the first three terms of this filtration.

The π_1-primage $X := \pi_1^{-1}(\Delta_{OAB})$ of the triangle $\Delta_{OAB} \subset Q_1$ is (the closure of) a fundamental domain of the $\mathbb{D}_8$-action on $S^n \times S^n$. It is described by the inequalities $0 \leq y_1 \leq x_1 \leq 1$ and as a subset of $S^n \times S^n$ it is an orientable manifold with boundary. This manifold has corners and possibly singularities of high codimension, however the associated fundamental class $x \in H_{2n-1}(X, \partial X)$ is well defined. The geometric boundary of X is,

$$\partial X = \pi_1^{-1}(\partial \Delta_{OAB}) = X_0' \cup X_1' \cup X_2' = \pi_1^{-1}(OA) \cup \pi_1^{-1}(OB) \cup \pi_1^{-1}(AB).$$

Homologically significant are only X_0' and X_1' since X_2' has codimension n in $S^n \times S^n$. Since $\mathrm{Stab}(X_0') = \langle \beta \rangle$ we subdivide and define $X_0 = X_0' \cap \{y_2 \geq 0\}$ as the associated fundamental domain. Similarly, $\mathrm{Stab}(X_1') = \langle \gamma \rangle$ and $X_1 = X_1' \cap \{y_2 \leq x_2\}$, cf. Figure 3 and the subsequent diagram.



(a) (b) (c)

Figure 3. Admissible filtration for $S^n \times S^n$.



We continue the "take the boundary, then subdivide"-procedure focusing as before only on the homologically significant part of the boundary.

$$\begin{array}{rclcl} \partial X_0 & = & Z_0' \cup T_0' & = & \{x_1 = y_1 = 0, y_2 \geq 0\} \cup \{x_1 \geq 0, y_1 = y_2 = 0\} \\ \partial X_1 & = & Z_0'' \cup T_1' & = & \{x_1 = y_1 = 0, y_2 \leq x_2\} \cup \{x_1 = y_1, x_1 \geq 0, x_2 = y_2\} \end{array}$$

The sets Z_0' and Z_0'' can be further subdivided as follows (Figure 3)

$$\begin{array}{rcl} Z_0' & = & Z_0 \cup \alpha Z_0 \cup \gamma Z_0 \cup \alpha\gamma Z_0 \\ Z_0'' & = & Z_0 \cup \beta Z_0 \cup \beta\gamma Z_0 \cup \alpha\beta\gamma Z_0 \end{array}$$

where $Z_0 := \{x_1 = y_1 = 0, 0 \leq y_2 \leq x_2 \leq 1\}$. Finally, $T_0' = T_0 \cup \beta T_0$ and $T_1' = T_1 \cup \gamma T_1$ where $T_0 = T_0' \cap \{0 \leq y_3 \leq 1\}$ and $T_1 = T_1' \cap \{y_3 \leq x_3\}$.

Summarizing we observe that the "take the boundary, then subdivide"-procedure defines an admissible filtration

$$S^n \times S^n = F_{2n} \supset F_{2n-1} \supset F_{2n-2} \supset \dots \tag{6.1}$$

where $F_{2n-1} := \bigcup_{g \in \mathbb{D}_8} g(\partial X)$ and $F_{2n-2} := \bigcup_{g \in \mathbb{D}_8} g(Z_0 \cup T_0 \cup T_1 \cup X_2)$.

7. Fragment of the chain complex for $S^n \times S^n$

The sets $X, X_0, X_1, Z_0, T_0, T_1$ described in the previous section are connected manifolds with boundary (with corners and possibly with mild singularities in codimension ≥ 2). They all can be oriented in which case the corresponding fundamental classes are denoted by $x, x_0, x_1, z_0, t_0, t_1$. These classes are naturally interpreted as the generators of $\mathbb{D}_8$-modules $H_k(F_k, F_{k-1}; \mathbb{Z})$ for $k = 2n, 2n-1, 2n-2$.

The orientation character of the $\mathbb{D}_8$-manifold $S^n \times S^n$ is given by

$$(\alpha, \beta, \gamma) = ((-1)^{n-1}, (-1)^{n-1}, (-1)^n). \tag{7.1}$$

From (7.1) and the analysis of geometric boundaries given in Section 6 one deduces the following relations:

$$\begin{array}{rcl} \partial x & = & (1 + (-1)^n\beta)x_0 + (1 + (-1)^{n-1}\gamma)x_1 \\ \partial x_0 & = & (1 + (-1)^n\alpha + (-1)^{n-1}\gamma - \alpha\gamma)z_0 + (1 + (-1)^{n-1}\beta)t_0 \\ \partial x_1 & = & -(1 + (-1)^n\beta - \beta\gamma + (-1)^{n-1}\alpha\beta\gamma)z_0 + (1 + (-1)^n\gamma)t_1 \end{array} \tag{7.2}$$

The top dimensional fragment of the associated chain complex is,

$$\Lambda x \xrightarrow{B} \Lambda x_0 \oplus \Lambda x_1 \xrightarrow{A} \Lambda z_0 \oplus \Lambda t_0 \oplus \Lambda t_1 \longrightarrow \cdots \tag{7.3}$$

The boundary homomorphisms is described by (7.2) so for example if n is even,

$$A^t = \left[\begin{array}{ccc} 1 + \alpha - \gamma - \alpha\gamma & 1 - \beta & 0 \\ -(1 + \beta - \beta\gamma - \alpha\beta\gamma) & 0 & 1 + \gamma \end{array} \right] \quad B = \left[\begin{array}{c} 1 + \beta \\ 1 - \gamma \end{array} \right] \tag{7.4}$$

8. Evaluation of the cohomology group $\mathcal{H}^{2n-1}_{\mathbb{D}_8}(S^n \times S^n; \mathcal{Z})$

The first obstruction to the existence of an equivariant map (5.2) is evaluated in the group $\mathcal{H}^{2n-1}_{\mathbb{D}_8}(S^n \times S^n; \mathcal{Z})$ where $\mathcal{Z} \cong H_{3j-1}(S(W^{\oplus j}); \mathbb{Z})$ is the orientation character of the sphere $S(W^{\oplus j})$. We remind the reader that these groups are defined (Section 11) as functors of $\mathbb{D}_8$-spaces with admissible filtrations. Also note that the condition $2d - 3j = 1$ (Section 2) allows us to assume that j is an odd integer.

Proposition 8.1. *Let $\mathcal{M} = \mathcal{Z}$ be the orientation character of the sphere $S(W^{\oplus j})$ where j is an odd integer. Then,*

$$\mathcal{H}^{2n-1}_{\mathbb{D}_8}(S^n \times S^n; \mathcal{Z}) \cong \left\{ \begin{array}{ll} \mathbb{Z}/4 & \text{if } n \text{ is even;} \\ \mathbb{Z}/2 \oplus \mathbb{Z}/2 & \text{if } n \text{ is odd.} \end{array} \right. \tag{8.2}$$

Moreover a generator of $\mathbb{Z}/4$ can be described as the cocycle

$$\phi : \Lambda x_0 \oplus \Lambda x_1 \to \mathcal{Z}, \quad \phi(x_0) = 1, \phi(x_1) = -1. \tag{8.3}$$

Proof. The orientation character $\mathcal{Z}$ of the $\mathbb{D}_8$-sphere $S(W)$ is easily determined from the signs of permutations (4.7) and reads as follows

$$(\alpha, \beta, \gamma) = (+1, +1, -1).$$

Since j is an odd integer the same answer is obtained for the orientation character of the sphere $S(W^{\oplus j})$. Assume that n is even. By applying the functor $\mathrm{Hom}(\cdot, \mathcal{Z})$ on the chain complex (7.3) we obtain the complex

$$\mathbb{Z} \xleftarrow{B_1} \mathbb{Z} \oplus \mathbb{Z} \xleftarrow{A_1} \mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z} \tag{8.4}$$

where,

$$A_1 = \left[\begin{array}{ccc} 4 & 0 & 0 \\ -4 & 0 & 0 \end{array} \right] \quad B_1^t = \left[\begin{array}{c} 2 \\ 2 \end{array} \right]$$

From here we deduce that

$$\mathcal{H}^{2n-1}_{\mathbb{D}_8}(S^n \times S^n; \mathcal{Z}) \cong \mathbb{Z}/4 \tag{8.5}$$

where a generator is the cocycle $\phi : \Lambda x_0 \oplus \Lambda x_1 \to \mathcal{Z}$, $\phi(x_0) = 1, \phi(x_1) = -1$. The second half of (8.2) is established by a similar calculation. □

9. Evaluation of the obstruction

In light of Propositions 5.1 and 8.1 (isomorphisms (8.2)) our primary concern are admissible triples $(d, j, 2)$ such that d is an even integer; this is precisely the case when the obstruction group is $\mathbb{Z}/4$. Hence, $(d, j) = (6k + 2, 4k + 1)$ for some integer $k \geq 0$. The following proposition is a key result for the evaluation of the primary obstruction $\theta \in \mathbb{Z}/4$ for the existence of a $\mathbb{D}_8$-equivariant map (5.2).

Proposition 9.1. *Suppose that*

$$\phi : S^{d-1} \times S^{d-1} \to S(W^{\oplus j}) \tag{9.2}$$

is a $\mathbb{D}_8$-equivariant map where $(d, j) = (6k + 2, 4k + 1)$ for some integer $k \geq 0$. Then,

$$\deg(\phi) \equiv 4\binom{2k-1}{k-1} \pmod 8. \tag{9.3}$$

We postpone the proof of Proposition 9.1 until we make some useful preliminary observations. It is not difficult to see that an equivariant map (9.2) must exist. However, we construct a particularly nice map which will make the evaluation of the degree easier.

Let $\mathcal{P}_m \cong \mathbb{R}^{m+1}$ be the vector space of all polynomials with real coefficients of degree $\leq m$, and $\mu_{m,n} : \mathcal{P}_m \times \mathcal{P}_n \to \mathcal{P}_{m+n}$, $\mu_{m,n}(p, q) = p \cdot q$, the multiplication of polynomials of indicated degrees. This map is non-degenerate in the sense that $\mu_{m,n}(p, q) = 0$ implies that either $p = 0$ or $q = 0$. As a consequence it induces a map

$$\mu'_{m,n} : \mathcal{P}'_m \times \mathcal{P}'_n \to \mathcal{P}'_{m+n}$$

where $\mathcal{P}'_m := \mathcal{P}_m \setminus \{0\}$ is the set of non-zero polynomials.

Proposition 9.4. *Suppose that $m = 2k$ and $n = 2l$ are even integers. The degree of the map $\mu'_{m,n} : \mathcal{P}'_m \times \mathcal{P}'_n \to \mathcal{P}'_{m+n}$ is*

$$\deg(\mu'_{m,n}) = \pm 2 \cdot \binom{k+l}{k}. \tag{9.5}$$

Proof. Let $\mathcal{P}^0_m := \{a_0 + \ldots + a_m t^m \in \mathcal{P}_m \mid a_m = 1\}$ be the hyperplane of monic polynomials. The radial projection is a bijection of $\mathcal{P}^0_m$ with the open hemisphere $S^n_+ := \{p \in \mathcal{P}_m \mid \|p\| = 1,\, a_m > 0\}$. Similarly $-\mathcal{P}^0_m$ is radially projected on the negative (open) hemisphere S^n_-. From here we conclude that the restriction

$$\mu^0 = \mu^0_{m,n} : \mathcal{P}^0_m \times \mathcal{P}^0_n \to \mathcal{P}^0_{m+n} \tag{9.6}$$

is a proper map and $\deg(\mu'_{m,n}) = 2 \cdot \deg(\mu^0)$. The degree $\deg(\mu^0)$ is computed by an algebraic count of the number of points in the pre-image $(\mu^0)^{-1}(\rho)$ of a carefully chosen generic polynomial $\rho \in \mathcal{P}^0_{m+n}$.

Assume that $\rho = \rho_1\rho_2 \dots \rho_{k+l}$ is a product of pairwise distinct, irreducible, quadratic (monic) polynomials ρ_i. Equivalently ρ does not have real roots and all its roots are pairwise distinct. Note that such a polynomial can be found in any neighborhood of the polynomial t^{m+n}. It follows that ρ is a regular value of μ^0 and

$$(\mu^0)^{-1}(\rho) = \{(p, q) \in \mathcal{P}^0_m \times \mathcal{P}^0_n \mid p \cdot q = \rho\}.$$

It is not difficult to see that all these points contribute to the degree with the same sign. Indeed, the determinant of the linear map $d\mu^0_{(p,q)}$, in an appropriate polynomial basis, is essentially the resultant $R(p, q)$ of polynomials p and q. It follows that

$$\deg(\mu^0_{m,n}) = \pm\binom{k+l}{k}$$

which implies the formula (9.5). □

Corollary 9.7. *Suppose that $f : S^m \times S^m \to S(\epsilon^{\oplus(2m+1)})$ is a $\mathbb{D}_8$-equivariant map where $m = 2k$ is an even integer. Then*

$$\deg(f) \equiv 4\binom{2k-1}{k-1} \pmod 8.$$

Proof. It is a well known fact, see [2, Section II.4], [12] or [23, Proposition 11], that under fairly general conditions G-equivariant maps have degrees congruent modulo $|G|$. In light of Proposition 9.4 it is sufficient to observe that the multiplication map $\mu_{m,m} : \mathcal{P}_m \times \mathcal{P}_m \to \mathcal{P}_{2m}$ is $\mathbb{D}_8$-equivariant and $\mathcal{P}_{2m} \cong \epsilon^{\oplus(2m+1)}$ as $\mathbb{D}_8$-modules. □

Example 9.8. The $\mathbb{D}_8$-equivariant map $\mu : S^2 \times S^2 \to S^4$ associated to the multiplication of quadratic polynomials $\mu'_{2,2} : \mathcal{P}'_2 \times \mathcal{P}'_2 \to \mathcal{P}'_4$ has degree ±4, consequently the degree of any $\mathbb{D}_8$-equivariant map is an integer of the form $8k + 4$.

Proof of Proposition 9.1. As in the proof of Corollary 9.7 it is sufficient to exhibit a particular $\mathbb{D}_8$-equivariant map (9.2) which satisfies the congruence (9.3).

Let us construct a $\mathbb{D}_8$-equivariant map

$$\Phi : \mathbb{R}^d \times \mathbb{R}^d \to W^{\oplus j} \tag{9.9}$$

with the property that if $\Phi(p, q) = 0$ then either $p = 0$ or $q = 0$. By decomposing the real $\mathbb{D}_8$-modules $\mathbb{R}^d \times \mathbb{R}^d \cong U^{\oplus d} \cong U^{\oplus j} \oplus U^{\oplus(d-j)}$ and $W^{\oplus j} \cong U^{\oplus j} \oplus \epsilon^{\oplus j}$ (Section 4.2) we observe that it is sufficient to construct a $\mathbb{D}_8$-equivariant map

$$\Phi' : \mathbb{R}^{d-j} \times \mathbb{R}^{d-j} \to \epsilon^{\oplus j} \tag{9.10}$$

which is non-degenerate in the sense that if $\Phi'(p, q) = 0$ then either $p = 0$ or $q = 0$. Let $m := 2k$ so $(d, j) = (3m + 2, 2m + 1) = (6k + 2, 4k + 1)$. Identify $\mathbb{R}^{d-j}$ with the space of real polynomials of degree less or equal to $d - j - 1 = m$ and $\epsilon^{\oplus j}$ with the space of real polynomials of degree $\leq j - 1 = 2m$. Then the multiplication of polynomials defines a non-degenerate (symmetric) bilinear form $\Phi'(p, q) = p \cdot q$ which is an example of a $\mathbb{D}_8$-equivariant map (9.10) with the desired properties.

Summarizing, using the identifications $\mathbb{R}^d \cong \mathcal{P}_{2m} \oplus \mathcal{P}_m$ and $W^{\oplus j} \cong (\mathcal{P}_{2m})^{\oplus 3}$ with the corresponding vector spaces of polynomials, we observe that the map Φ has the following explicit form,

$$\Phi(p, q) = \Phi(p', p''; q', q'') = (p', q', p''q'') \tag{9.11}$$

where $p''q'' = \mu(p'', q'')$ is the polynomial multiplication. The degree $\deg(\Phi)$ can be calculated again, as in the proof of Proposition 9.4, by the reduction to the case of monic polynomials p'' and q''. By choosing the regular value of the map (9.11) in the form $(0, 0, \rho)$, where ρ is a regular value for the multiplication of monic polynomials we observe that Proposition 9.1 is an immediate consequence of Proposition 9.4. □

The following proposition is the central result of this section and the ultimate goal of all earlier computations. Note that the manifold $S^{d-1} \times S^{d-1}$ (Proposition 9.1) can be naturally identified as a subset of the $(2d-2)$-dimensional element F_{2d-2} of the filtration (6.1) via the identification $S^{d-1} \times S^{d-1} = (S^d \times S^d) \cap \{x_1 = y_1 = 0\} = \bigcup_{g \in \mathbb{D}_8} g(Z_0)$.

Proposition 9.12. *Suppose that* $\Delta = 2d - 3j = 1$ *where* $(d, j) = (3m+2, 2m+1)$ *and* $m = 2k$ *is an even integer. Then the first obstruction to the existence of a* $\mathbb{D}_8$*-equivariant map (5.2), evaluated as an element of* $\mathbb{Z}/4$*, is equal to*

$$\theta = 2\binom{2k-1}{k-1} \pmod 4. \tag{9.13}$$

Proof. By (8.3) a generator of the obstruction group is the cocycle ϕ such that $\phi(x_0) = 1, \phi(x_1) = -1$ where x_0 and x_1 are the (relative) fundamental classes of the pseudomanifolds X_0 and X_1. Following the calculations (and notation) from Section 6, the geometric boundary of the pseudomanifold X_0 has the following representation.

$$\partial X_0 = Z_0' \cup T_0' = (Z_0 \cup \alpha Z_0 \cup \gamma Z_0 \cup \alpha\gamma Z_0) \cup (T_0 \cup \beta T_0). \tag{9.14}$$

The set $M := Z_0' \cup T_0'$ is, up to a closed subset of high codimension, a closed oriented manifold. In light of (11.14) the obstruction θ can be evaluated as the degree $\deg(f)$ where $f = \psi|_M$ is the restriction on M of an arbitrary $\mathbb{D}_8$-equivariant map $\psi : F_{2d-2} \to S(U^{\oplus j} \oplus \epsilon^{\oplus j})$. Such a map clearly exists since $\dim(F_{2d-2}) = \dim(S(W^{\oplus j}))$. Interpreting the degree as an algebraic count of points in the preimage of a regular value, we observe that

$$\deg(\psi) = \deg(\psi_{|Z_0'}) + \deg(\psi_{|T_0'}).$$

Using the $\mathbb{D}_8$-symmetries of X_0 and T_0, in particular the fact that X_0 is "a half of the manifold" $S^{d-1} \times S^{d-1} \subset F_{2d-2}$, together with Proposition 9.1, we observe that

$$\deg(\psi_{|Z_0'}) = 2\binom{2k-1}{k-1} \quad \text{and} \quad \deg(\psi_{|T_0'}) = 0$$

which completes the proof of the proposition. □

10. SUMMARY OF PROOFS OF MAIN RESULTS

Proof of Theorem 2.1. By Proposition 5.1 Theorem 2.1 is an immediate consequence of Theorem 2.3. □

Proof of Theorem 2.3. By Proposition 9.12 the first obstruction to the existence of an equivariant map (2.4) is given by (9.13). It is non-zero if and only if $\binom{2k-1}{k-1}$

is an odd integer which is the case if and only if $k = 2^l$. It follows that in the case when d is even the triple $(d, j, 2)$ is admissible if for some integer $l \geq 0$,

$$d = 3 \cdot 2^{l+1} + 2 \quad \text{and} \quad j = 2 \cdot 2^{l+1} + 1.$$

11. Appendix: obstructions, filtrations and chain complexes

For a review of equivariant obstruction theory the reader is referred to [2, Section II.3]. One of the central results is the following *obstruction exact sequence.*

Theorem 11.1. *Suppose that X is a free G-CW-complex and that Y is n-simple G-space. Then for each integer $n \geq 1$ there exists an exact sequence,*

$$[X^{(n+1)}, Y]_G \longrightarrow \mathrm{Image}\{[X^{(n)}, Y]_G \to [X^{(n-1)}, Y]_G\} \longrightarrow \mathcal{H}^{n+1}_G(X, \pi_n(Y)) \tag{11.2}$$

In many applications $Y = S(V)$ is a G-sphere $S(V) \cong S^n$ for some real G-module V. In that case Theorem 11.1 has the following important corollary.

Corollary 11.3. *Suppose that X is a free G-CW-complex. Let $Y = S(V) = S^n$ be an n-dimensional G-sphere associated to a real G-representation V $(n \geq 2)$. Then (11.2) reduces to,*

$$[X^{(n+1)}, S(V)]_G \longrightarrow \{*\} \longrightarrow \mathcal{H}^{n+1}_G(X, \mathcal{Z}) \tag{11.4}$$

where $\mathcal{Z}$ is the orientation character of $S(V)$ and $\{*\}$ is a singleton.

The exactness of the sequence (11.4) means that there exists a single element of the obstruction group $\mathcal{H}^{n+1}_G(X, \mathcal{Z})$ (the image of $*$) which is zero if and only if there exists a G-equivariant map $f : X^{(n+1)} \to S(V)$.

Our objective is to extend the applicability of these results by allowing more general filtrations which not necessarily arise from a G-CW-structure on X. The reader may keep on mind the G-manifold complexes introduced in Section 4.1 as a guiding example.

Definition 11.5. Let X be a not necessarily free G-space which admits a G-invariant triangulation (CW-structure) turning X into a simplicial complex (CW-complex) of dimension $m = n + 1$. Let $X^{(k)}$ be the associated k-skeleton. A finite filtration

$$\emptyset = X_{-1} \subset X_0 \subset X_1 \subset \ldots \subset X_{n-1} \subset X_n \subset X_{n+1} = X \tag{11.6}$$

is called *admissible* if,

(1) X_k is a G-invariant subcomplex of $X^{(k)}$ for each k.

Let $C_*(X)$ be the G-chain complex associated to the filtration (11.6) where $C_k(X) := H_k(X_k, X_{k-1}; \mathbb{Z})$. The associated cohomology groups with coefficients in a G-module M are denoted by

$$\mathcal{H}^*(X; M). \tag{11.7}$$

Remark 11.8. The reader should keep in mind that the cohomology groups (11.7) are functors of a filtered space X, not the space alone. This is in agreement with the approach in [2, page 112] where the corresponding equivariant cohomology groups depend on a given G-CW-structure.

An admissible filtration (11.6) is *free* if X is a free G-space and the associated chain complex is free, i.e.

(2) $C_k = C_k(X) := H_k(X_k, X_{k-1}; \mathbb{Z})$ is a free Λ-module, where $\Lambda := \mathbb{Z}[G]$ is the group ring of the group G.

Remark 11.9. The structure of a simplicial complex on a G-space X plays an auxiliary role and in intended applications one starts directly with a filtration (11.6), silently assuming that it can be "triangulated". The most natural is the situation when X is a G-manifold complex where the constituent manifolds are semialgebraic sets (as in Section 6) and this condition is automatically satisfied. The condition (2) is evidently not necessarily satisfied if X is a free G-space with an admissible filtration.

We assume that the target G-space Y is also filtered by a filtration

$$\emptyset = Y_{-1} \subset Y_0 \subset Y_1 \subset \ldots \subset Y_n \subset Y_{n+1} \subset \ldots \subset Y_\nu = Y \tag{11.10}$$

arising from some, not necessarily free, G-CW-structure on Y. Let $D_* = (\{D_k\}_0^\nu, \partial)$ be the associated cellular chain complex, $D_k := H_k(Y_k, Y_{k-1})$.

The following proposition allows us to reduce the problem of the existence of G-equivariant maps between X and Y, to the question of the existence of chain maps between the associated chain complexes of $\Lambda = \mathbb{Z}[G]$ modules.

Proposition 11.11. *Suppose that X is an $(n+1)$-dimensional G-space with an* admissible filtration *(11.6) (Definition 11.5). Suppose that Y is a G-CW-complex and let (11.10) be the associated filtration by skeletons. Then if there exists a G-equivariant map $f : X \to Y$, there exists also a chain map $f_* : C_*(X) \to D_*(Y)$ of the associated, augmented chain complexes,*

(11.12)

$$\begin{tikzcd}
C_{n+1} \arrow[r, "\partial"] \arrow[d, "f_{n+1}"] & C_n \arrow[r, "\partial"] \arrow[d, "f_n"] & C_{n-1} \arrow[r] \arrow[d, "f_{n-1}"] & \cdots \arrow[r] & C_1 \arrow[r, "\partial"] \arrow[d, "f_1"] & C_0 \arrow[r] \arrow[d, "f_0"] & \mathbb{Z} \arrow[r] \arrow[d, "\cong"] & 0 \\
D_{n+1} \arrow[r] & D_n \arrow[r] & D_{n-1} \arrow[r] & \cdots \arrow[r] & D_1 \arrow[r] & D_0 \arrow[r] & \mathbb{Z} \arrow[r] & 0
\end{tikzcd}$$

Proof. If there exists a G-equivariant map $f : X \to Y$ then by the *cellular approximation theorem* (Theorem 2.1 in [2, Section II.2]) there exists a cellular map $g : X \to Y$ which is G-homotopic to f. In follows that $g(X_k) \subset g(X^{(k)}) \subset Y^{(k)} = Y_k$ which in turn implies the existence of the chain map (11.12). □

Proposition 11.11 allows us to reduce the topological problem of the existence of equivariant maps to an algebraic problem of finding a chain map. This in turn leads to algebraic counterparts of Theorem 11.1 and Corollary 11.3.

Proposition 11.13. *Suppose that $C_* := \{C_k\}_{k=-1}^{n+1}$ and $D_* := \{D_k\}_{k=-1}^{n+1}$ are chain complexes of $\Lambda = \mathbb{Z}[G]$ modules where $C_{-1} \cong D_{-1} \cong \mathbb{Z}$. Suppose that the chain map $F_{n-1} := (f_j)_{j=-1}^{n-1} : \{C_k\}_{k=-1}^{n-1} \to \{D_k\}_{k=-1}^{n-1}$ exists and is fixed in advance. Suppose that F_{n-1} can be extended one step further i.e. that there exists a homomorphism $\hat{f}_n : C_n \to D_n$ such that $\partial \circ \hat{f}_n = f_{n-1} \circ \partial$. Then the obstruction to the existence of a chain map (11.12), $F_{n+1} := (f_j)_{j=-1}^{n+1} : \{C_k\}_{k=-1}^{n+1} \to \{D_k\}_{k=-1}^{n+1}$, extending the chain map F_{n-1}, is a well defined element θ of the cohomology group*

$$H^{n+1}(C_*; H_n(D_*)) = H_{n+1}(\mathrm{Hom}(C_*, H_n(D_*))).$$

Moreover, θ is represented by the cocycle

$$\theta(\hat{f}_n) : C_{n+1} \xrightarrow{\partial} C_n \xrightarrow{\hat{f}_n} Z_n(D) \xrightarrow{\pi} H_n(D). \tag{11.14}$$

The vanishing of θ is not only necessary but also sufficient for the existence of the chain map (11.12) if the modules C_n and C_{n+1} are free (projective).

Proof. The proof is routine and left to the reader. ☐

Remark 11.15. The last part of Proposition 11.13, claiming that θ is a complete obstruction provided C_n and C_{n+1} are free modules, is the main motivation for working with free admissible filtrations, cf. condition (2) in Definition 11.5. Note that the freeness of C_n and C_{n+1} is a condition that can be satisfied even if X is not a free G-space, e.g. if the corresponding set of fixed points X^H has a codimension ≥ 2 for each subgroup $H \neq \{e\}$.

Acknowledgments. It is a pleasure to thank P. Landweber, S. Melikhov, and members of the Belgrade CGTA[2] seminar for useful comments on a preliminary version of this paper (April 10, 2011).

References

[1] P.V.M. Blagojević and G.M. Ziegler. The ideal-valued index for a dihedral group action, and mass partition by two hyperplanes, arXiv:0704.1943v2 [math.AT]), 9 Dec. 2010.

[2] T. tom Dieck. *Transformation Groups.* de Gruyter Stud. Math., vol. 8, Springer, Berlin, 1987.

[3] J. González and P. Landweber. The integral cohomology groups of configuration spaces of pairs of points in real projective spaces, arXiv:1004.0746v2 [math.AT], 17 Dec 2010.

[4] M. Gromov. Isoperimetry of waists and concentration of maps. *GAFA, Geom. Funct. Anal.*, 13 (2003), 178–215.

[5] B. Grünbaum. Partitions of mass–distributions and convex bodies by hyperplanes. *Pacific J. Math.*, 10 (1960), 1257–1261.

[6] L. Guth. The endpoint case of the Bennett-Carbery-Tao multilinear Kakeya conjecture, arXiv:1103.2926v2 [math.CO].

[7] L. Guth and N.H. Katz. On the Erdös distinct distance problem in the plane, arXiv:1011.4105v1 [math.CO].

[8] H. Hadwiger. Simultane Vierteilung zweier Körper. *Arch. Math.* (Basel), 17 (1966), 274–278.

[9] A. Hubard and B. Aronov. Convex Equipartitions of volume and surface area, arXiv:1010.4611v3 [math.MG].

[10] R.N. Karasev. Equipartition of several measures, arXiv:1011.4762v6 [math.MG].

[11] M. Kreck. *Differential Algebraic Topology.* Graduate Studies in Mathematics, vol. 110, American Mathematical Society 2010.

[12] A. Kushkuley and Z.I. Balanov. *Geometric Methods in Degree Theory for Equivariant Maps.* Lecture Notes in Math. 1632, Springer, Berlin 1996.

[13] P. Mani-Levitska, S. Vrećica, and R. Živaljević. Topology and combinatorics of partitions of masses by hyperplanes. *Adv. Math.*, 207 (2006), 266-296. MR 2264074

[14] J. Matoušek. The dawn of an algebraic era in Discrete Geometry? `http://kam.mff.cuni.cz/~matousek/mor.pdf`.

[15] B. Matschke. A note on masspartitions by hyperplanes, arXiv:1001.0193v1 [math.CO], 31 Dec 2009.

[16] E.A. Ramos. Equipartitions of mass distributions by hyperplanes. *Discrete Comput. Geom.*, vol. 15 (1996), pp. 147–167.

---

[2] Combinatorics in Geometry, Topology, and Algebra.

[17] M. Sharir. From joints to distinct distances and beyond: The dawn of an algebraic era in combinatorial geometry, `http://www.cs.tau.ac.il/~michas/algober.pdf`.
[18] P. Soberón. Balanced convex partitions of measures in $\mathbb{R}^d$, arXiv:1010.6191v2 [math.MG].
[19] J. Solymosi and T. Tao. An incidence theorem in higher dimensions, arXiv:1103.2926v2 [math.CO].
[20] H. Steinhaus. A note on the ham sandwich theorem. *Mathesis Polska* 9 (1938), 26-28.
[21] A.H. Stone, J.W. Tukey. Generalized sandwich theorems. *Duke Math. J.* 9 (1942), 356-359.
[22] T. Tao. `http://terrytao.wordpress.com/?s=polynomial+ham+sandwich`
[23] S.T. Vrećica, R.T. Živaljević. Chessboard complexes indomitable, arXiv:0911.3512v3 [math.CO], 10 Apr 2011.
[24] R. Živaljević. The Tverberg-Vrećica problem and the combinatorial geometry on vector bundles. *Israel J. Math*, 111:53-76, 1999.
[25] R.T. Živaljević. Topological methods. Chapter 14 in *Handbook of Discrete and Computational Geometry*, J.E. Goodman, J. O'Rourke, eds, Chapman & Hall/CRC 2004, 305–330.
[26] R.T. Živaljević. Equipartitions of measures in $\mathbb{R}^4$. *Trans. Amer. Math. Soc.* 360 (2008), 153–169.

Mathematical Institute SASA, Knez Mihailova 36, 11001 Belgrade, Serbia
E-mail address: `rade@turing.mi.sanu.ac.rs`